

Europäisches Patentamt

European Patent Office

Office européen des brevets



(11) EP 1 048 295 A2

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication: 02.11.2000 Bulletin 2000/44

(21) Application number: 00303077.2

(22) Date of filing: 12.04.2000

(51) Int. CI.⁷: **A61K 31/216**, A61P 3/06, A61K 9/14, A61K 9/48

Willowdale, Ontario M2L 2K1 (CA)

(84) Designated Contracting States:

AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU MC NL PT SE Designated Extension States: AL LT LV MK RO SI

(30) Priority: 27.04.1999 CA 2270306

(71) Applicant: SHERMAN, Bernard Charles Willowdale, Ontario M2L 2K1 (CA) (72) Inventor: SHERMAN, Bernard Charles

(74) Representative:
Votier, Sidney David
CARPMAELS & RANSFORD
43, Bloomsbury Square
London WC1A 2RA (GB)

(54) Pharmaceutical compositions comprising co-micronized fenofibrate

(57) A pharmaceutical composition for oral administration comprising a co-micronized mixture of fenofibrate and a solid excipient that is not a surfactant.

FIELD OF INVENTION

[0001] The present invention relates to pharmaceutical compositions for oral administration comprising fenofibrate which enable improve dissolution and bioavailability.

1

BACKGROUND

[0002] Fenofibrate is practically insoluble in water. This causes fenofibrate to exhibit a low rate of dissolution in aqueous media (including gastrointestinal fluids). Which results in inadequate bioavailability (absorption into systemic circulation) after oral ingestion.

[0003] In order to make a composition comprising fenofibrate that will enable maximum bioavailability, it is necessary to incorporate into the composition a feature that increases the rate of dissolution of the drug in gastrointestinal fluids.

[0004] Several ways of increasing the rate of dissolution of drugs having low solubility in water are known in the prior art.

[0005] One approach is micronization. In this 25 approach, the drug is milled to fine particles, typically having a mean diameter of under about 15 microns. A second approach is to include a surfactant in the composition.

[0006] For the drug fenofibrate, neither micronization alone nor use of a surfactant alone enables maximum bioavailability. US Patent 4895726 discloses that the rate of dissolution and the bioavailability of fenofibrate can be maximized by co-micronization of fenofibrate with a solid surfactant. In this process the fenofibrate is first mixed with the solid surfactant and then the mixture is micronized.

[0007] A composition made according to the invention of US Patent 4895726 is sold in Canada under the tradename Lipidil Micro and in the United States under the tradename Tricor.

A disadvantage of the technology of US Patent 4895726 is the need to include the solid surfactant in the composition. Because of the toxicity of surfactants, it is preferable to avoid use of a surfactant if possible.

Another method of increasing the dissolution rate of fenofibrate is disclosed in Canadian patent application No. 2214895. This publication discloses that the bioavailability of fenofibrate can be improved by making a solid dispersion of a disintegrant in the fenofibrate. This is done by melting the fenofibrate, blending the disintegrant into the molten fenofibrate, and resolidifying the mixture. The resulting solid can then be ground up into granules and the granules used to make the final composition. For example, the granules can be filled into two-piece hard gelatin capsules.

[0008] A disadvantage of the method of Canadian patent application No. 2214895 is that it requires the

use of specialized equipment to make the molten blend.

[0009] In view of the limitations of the prior art, it is the object of the present invention to enable increased dissolution rate of fenofibrate without the need to incorporate a surfactant in the composition, and without the need to make a molten blend.

DESCRIPTION OF THE INVENTION

[0010] It has been found that the dissolution rate of fenofibrate can be substantially increased by co-micronization of fenofibrate with a pharmaceutically acceptable excipient that is not a surfactant. This is surprising in light of the US Patent 4895726 which teaches co-micronization only with a solid surfactant.

[0011] The term "pharmaceutically acceptable excipient" will be understood to mean any ingredient having no therapeutic activity and being nontoxic and thus suitable as an excipient.

[0012] Suitable excipients will include any of the excipients commonly used in pharmaceutical products, such as, for example, microcrystalline cellulose, lactose and starch, provided that such excipient is solid at room temperature and not a surfactant.

[0013] The ratio of the weight of the excipient to the weight to fenofibrate may be anywhere from about 1:100 to about 2:1, will preferably be from about 1:10 to about 3:2, and will most preferably be about 1:1.

[0014] The co-micronization of the fenofibrate and excipient will advantageously be carried out by mixing the fenofibrate and excipient together and then micronizing of the mixture on conventional micronization equipment, such as an air-jet mill. The mixture will preferably be micronized such that the mean particle size is less than 15 microns, more preferably less than 10 microns, and most preferably less than 5 microns.

[0015] The co-micronized powder may then be processed into solid dosage forms for oral administration (i.e. tablets or capsules).

[0016] This may be, for example, in one of the following ways:

- 1. Filling the co-micronized powder directly into 2piece hard gelatin capsules.
- 2. Mixing the co-micronized powder with other excipients, such as, for example, fillers, binders, disintegrants, lubricants and glidants, and either filling the mixture into 2-piece hard gelatin capsules or compressing the mixture into tablets.

[0017] The invention will be more clearly understood from the following examples.

Example 1

[0018] 500 g of fenofibrate was mixed with 500 g of lactose monohydrate powder, and the mixture was

45

15

25

4

micronized on an air-jet mill. 2 piece hard gelatin capsules were then filled with the resultant co-micronized powder to a net fill weight of 400 mg per capsule, so that each capsule contained 200 mg of fenofibrate.

Example 2

[0019] 500 g of fenofibrate was mixed with 500 g of microcrystalline cellulose, and the mixture was micronized on an air-jet mill. 2-piece hard gelatin capsules were then filled with the resultant co-micronized powder to a net fill weight of 400 mg per capsule, so that each capsule contained 200 mg of fenofibrate.

Example 3

[0020] For comparison purposes, a quantity of pure fenofibrate was micronized using the same air-jet mill.
[0021] A sample of the pure micronized fenofibrate was then mixed with an equal weight of lactose monohydrate power. 2-piece hard gelatin capsules were then filled with the resultant mixture to a net fill weight of 400 mg per capsule, so that each capsule again contained 200 mg of fenofibrate.

Dissolution Results

[0022] Capsules of examples 1 and 2 were compared to capsules of example 3 for dissolution rate.

[0023] The equipment used for dissolution testing 30 was United States Pharmacopoeia Apparatus #2. The paddle speed was 100 rpm, and the medium was 900 mL of 0.1N sodium dodecyl sulfate water.

[0024] It was found that, in 60 minutes, over 90% was dissolved from the capsules of examples 1 and 2, 35 whereas only 50% to 70% was dissolved for the capsules of example 3.

[0025] It is thus clear that the dissolution rate is substantially higher using fenofibrate that has been comicronized with a solid excipient such as lactose or 40 microcrystalline cellulose, in comparison to fenofibrate that has been micronized in pure form and then mixed with a solid excipient.

Claims

- A pharmaceutical composition comprising a comicronized mixture of fenofibrate and a solid excipient that is not a surfactant.
- A composition as in claim 1 wherein the mean particle size of the said co-micronized mixture is less than 15 microns.
- A composition as in claim 1 wherein the mean particle size of the said co-micronized mixture is less than 10 microns.

- A composition as in claim 1 wherein the mean particle size of the said co-micronized mixture is less than 5 microns.
- 5 5. A composition as in any of claims 1 to 4, wherein the ratio of the excipient to fenofibrate by weight is from about 1:100 to about 2:1.
 - A composition as in any of claims 1 to 4, wherein the ratio of the excipient to fenofibrate by weight is about 1:10 to about 3:2.
 - A composition as in any of claims 1 to 4, wherein the ratio of the excipient to fenofibrate by weight is about 1:1.
 - 8. A composition as in any of claims 1 to 5, wherein the excipient is selected from the group consisting of microcrystalline cellulose, lactose, and starch.

3

45



Europäisches Patentamt

European Patent Office

Office européen des brevets



EP 1 048 295 A3

(12)

EUROPEAN PATENT APPLICATION

- (88) Date of publication A3: 06.12.2000 Bulletin 2000/49
- (43) Date of publication A2: 02.11.2000 Bulletin 2000/44
- (21) Application number: 00303077.2
- (22) Date of filing: 12.04.2000

(51) Int. CI.⁷: **A61K 31/216**, A61P 3/06, A61K 9/14, A61K 9/48

(11)

(84) Designated Contracting States:

AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU

MC NL PT SE
Designated Extension States:
AL LT LV MK RO SI

- (30) Priority: 27.04.1999 CA 2270306
- (71) Applicant:
 SHERMAN, Bernard Charles
 Willowdale, Ontario M2L 2K1 (CA)

(72) Inventor:

SHERMAN, Bernard Charles
Willowdale, Ontario M2L 2K1 (CA)

- (74) Representative:
 Votier, Sidney David
 CARPMAELS & RANSFORD
 43, Bloomsbury Square
 London WC1A 2RA (GB)
- (54) Pharmaceutical compositions comprising co-micronized fenofibrate
- (57) A pharmaceutical composition for oral administration comprising a co-micronized mixture of fenofibrate and a solid excipient that is not a surfactant.



EUROPEAN SEARCH REPORT

Application Number EP 00 30 3077

	~ 	ERED TO BE RELEVANT		
Category	Citation of document with in of relevant pass	ndication, where appropriate, ages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.CI.7)
A,D	US 4 895 726 A (B.C 23 January 1990 (19 * claims *		1-8	A61K31/216 A61P3/06 A61K9/14 A61K9/48
A,D	EP 0 904 781 A (B.C 31 March 1999 (1999 * claims *		1-8	,
A	EP 0 724 877 A (LAB 7 August 1996 (1996 * claims *	ORATOIRES FOURNIER) -08-07)	1-8	
A	WO 96 21439 A (GALE 18 July 1996 (1996- * claims *		1-8	
				TECHNICAL FIELDS SEARCHED (Int.Cl.7)
				A61K
			i	
		·		
	The present search report has	been drawn up for all claims		
	Place of search	Date of completion of the search	ì	Examiner
	THE HAGUE	17 October 200	0 Sca	arponi, U
X : par Y : par doc	ATEGORY OF CITED DOCUMENTS toularly relevant if taken alone toularly relevant if combined with anounce of the same category.	E : earlier pater after the filln ther D : document d L : document d	ted in the application led for other reasons	Bshed on, or
O : nor	hnological background n-written disclosure amediate document	& : member of to document	he same palent fami	ly, corresponding

ANNEX TO THE EUROPEAN SEARCH REPORT ON EUROPEAN PATENT APPLICATION NO.

EP 00 30 3077

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

17-10-2000

(s)	Publication date
696 A	01-09-19
	15-01-19
	05-09-19
	31-08-19
	28-09-19
	28-01-19
	09-06-19
	30-08-19
	01-08-19
	30-06-19
	11-10-19
	25-10-19
	22-02-19
130 A	25-10-19
895 A	27-11-19
	15-04-19
	08-06-19
	09-08-19
	15-07-20
	03-08-20
	01-10-19
148 A	09-03-19
628 A	13-08-19
896 A	31-07-19
985 A	18-07-19
562 A	22-10-19
959 T	17-11-19

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82